

Inclusion criteria:

1. Patients ≥ 18 and < 80 years.
2. **Karnofsky** performance status previous to AP 100 (normal, no complaints, no evidence of disease), 90 (able to carry on normal activity; minor signs or symptoms of disease) or 80 (normal activity with effort; some signs or symptoms of disease).
3. Acute pancreatitis defined as at least 2 of the following 3 criteria: A) Typical abdominal pain, B) Elevation of amylase and/or lipase more than 3 times the upper limit of normality, and C) Imaging (preferably CT and/or MR) compatible with AP.
4. Written informed consent.

Exclusion criteria:

1. Time between onset of symptoms and presentation in the emergency room (ER) greater than **48 hours** (we do not consider as onset of symptoms previous self-limited heraldic biliary pain)
2. Recruitment > 24 h after presentation in the ER
3. Karnofsky performance status previous to AP 70 (Cares for self; unable to carry on normal activity or to do active work) or less (in general, patients who are sick before AP, so scoring in QoL scale will be low regardless of the severity of AP)
4. Inability to express or understand the instructions of the study (severe congenital or acquired intellectual deficit).
5. **More than 1 previous** episode of AP.
6. **Chronic pancreatitis.**
7. Presence of diseases or conditions different from AP that may interfere with the scale: other causes of abdominal pain (especially **acute cholecystitis**; NOTE: choledocholithiasis-acute cholangitis is not an exclusion criteria), obstruction of the digestive tract (peptic pyloric stenosis, gastrointestinal anastomotic stenosis, diabetic gastroparesis, gastrointestinal neoplasia...), nausea-vomiting (brain tumor, chemotherapy...) or weakness (pre-existing anemia with **Hb < 9 mg/dL**, heart failure or respiratory insufficiency associated with **minimal effort dyspnea**, or **domiciliary treatment with O₂**, advanced neoplasms or other **debilitating diseases**).

3. 11. 14

Hungarian

1. Fájdalom, különösen a has területén, mellkasi vagy háti fájdalom
2. Hasi feszülés (puffadás, fokozott gázképződés)
3. Korai teltségérzés, mintha az étel megakadna a gyomorban
4. Székürítési nehezítettség (székrekedés vagy erőlködő székelés)
5. Hányinger és/vagy hányás
6. Szomjúság
7. Gyengeség, fáradékonyság, erőtlenség, a mozgás nehezítettség

PROM Skála

24h	48h
(72h)	7.Nap ∅ 2p
5.Nap ∅ 2p	

Elbocsájtás napja

15. (±2) nap EORTC QLQ-C30 ∅ 4p